

## SOME INDICATORS OF DAMAGES AND RESPONSES OF *XANTHORIA PARIETINA* (L.) Th. Fr. TO FLUORIDE AND LEAD INDUCED-STRESS

 Ouahiba Benhamada<sup>1\*</sup>,  Nabila Benhamada<sup>2</sup>,  Essaid Leghouchi<sup>2</sup>

<sup>1</sup>University of Jijel, Faculty of Nature and Life Sciences, Laboratory of Biotechnology, Environment and Health, Department of Applied Microbiology and Food Sciences, Jijel, Algeria

<sup>2</sup>University of Jijel, Faculty of Nature and Life Sciences, Laboratory of Biotechnology, Environment and Health, Department of Cell and Molecular Biology, Jijel, Algeria

\*Corresponding Author:

E-mail: [wahibabenhamada@yahoo.fr](mailto:wahibabenhamada@yahoo.fr)

(Received 17<sup>th</sup> July 2022; accepted 13<sup>th</sup> October 2022)

**ABSTRACT.** One of the major problems of current time is air pollution, the assessment of air quality through the use of bioindicators is a major concern at the moment. Our work aims to study the response of *Xanthoria parietina* (L.) Th. Fr. to fluoride- and lead-induced stress as an example of the phytotoxic air pollutants. For this purpose, lichen thalli have been treated with sodium fluoride (NaF) and lead nitrates (Pb(NO<sub>3</sub>)<sub>2</sub>) solutions at 0, 0.5, 1.0, 5.0 and 10 mM, for time scale of 0, 24, 48 and 96 h. Lipid peroxidation measured by Malondialdehyde (MDA) and chlorophyll degradation measured by optical density OD<sub>435</sub>/OD<sub>415</sub> ratio are used as results of stress induced by fluoride and lead, and the accumulation of proline and soluble sugars are measured as indicators of responses used by *X. parietina*. Based on the obtained results, it was noted that lipid peroxidation increased correlating with increasing concentrations of NaF and Pb(NO<sub>3</sub>)<sub>2</sub> ( $r=0.773$ ,  $p=0.000712^{***}$  and  $r=0.865$ ,  $p=0.000031^{***}$ , respectively), however, chlorosis and proline increased correlating with increasing exposure time of NaF ( $r=-0.737$ ,  $p=0.0011^{**}$  and  $r=0.783$ ,  $p=0.00032^{***}$ , respectively) and Pb(NO<sub>3</sub>)<sub>2</sub> ( $r=-0.926$ ,  $p<0.0001^{***}$  and  $r=0.811$ ,  $p=0.00013^{***}$ , respectively), whereas soluble sugar contents increased according to increasing concentrations of NaF ( $r=0.678$ ,  $p=0.0010^{***}$ ) and according to increasing exposure time of Pb(NO<sub>3</sub>)<sub>2</sub> ( $r=0.780$ ,  $p=0.00036^{***}$ ). Although lead was significantly more toxic than fluoride ( $p=0.02^*$ ), *X. parietina* offers a very high sensitivity to fluoride, which allowed us to conclude that the toxicity of fluorine is comparable to that of lead.

**Keywords:** Chlorophyll, fluoride, lead, MDA, proline, soluble sugars.

### INTRODUCTION

Lichens present a very important model of symbiotic organisms associating a mushroom called mycobiont and green algae and/or cyanobacteria called photobiont [1], one of its main uses is the use as bioindicators of air quality [2, 3, 4]. Lichens have the ability to absorb significant amounts of trace elements from the atmosphere [5, 6]. Lichens are also used as biomonitors [7, 8, 9] and bioaccumulators of heavy metals [10, 11, 12, 13]. They are capable of growing on difficult supports such as coastal rocks [14], and differed from most other eukaryotic organisms in their physiology, morphology and

their ability to tolerate extreme stresses [15]. Despite their distribution and diversity are influenced by climate, soil chemistry and geography [16], lichens can resist various stressful conditions such as drought and temperature extremes [17], salinity [18], heat [19] heavy metals [12] and nutrient deficiency [20]. These stresses are the source of reactive oxygen species (ROS), the most important adaptation mechanism used by lichens for tolerance to stressful conditions is the scavenging of reactive oxygen species (ROS) [21]. Lichens can also respond to stressful situations by displaying conventional stress-tolerant traits including reduced growth rates, considerable longevity, low nutrient demand and the presence of particular morphological and physiological adaptation and changes in ecological behaviour for surviving in the world most hostile situations [22].

Stressful conditions caused several biochemical and physiological changes in the plants. The most commonly used parameters to study the toxicity of atmospheric pollutants on lichens are chlorophyll degradation [23, 24] and lipid peroxidation [25, 26].

Exposure to abiotic stress triggers the accumulation of amino acids and amines in different plant species. Soluble sugars play a crucial role in a variety of metabolic processes, acting as a signal to control gene expression in photosynthesis, osmolyte production, and sucrose metabolism [27]. Proline, on the other hand, is crucial for plants; it protects plants from various stresses and helps in their faster recovery from stress [28].

Our work which relates to the study of the toxic effect induced by fluoride and lead on *X. parietina*, is the first to be carried out in our region.

The aim of this study is to investigate the toxic effect of fluoride and lead on lipid peroxidation and chlorophyll degradation in *X. parietina* (L.) Th. Fr. lichen, and to check for the accumulation of proline and soluble sugars as indicators of responses used for adaptation to induced stress.

## MATERIALS AND METHODS

### *Lichen material*

Lichen thalli samples of *X. parietina* were gathered in the Beni Metrane and Djimar region, south of Jijel (Algeria) in February-March 2018. After collection, samples were transferred to the laboratory in clean closed boxes. Impurities were removed, and samples were washed with distilled water to remove superficial dust and adherent particles. In each experimental vessel, fresh weights of thalli were isolated and acclimatized to laboratory conditions until analysis.

### *Fluoride and lead treatment*

In comparison to distilled water, the lichen thalli of *X. parietina* were incubated in solutions of NaF and  $Pb(NO_3)_2$  at 0, 0.5, 1.0, 5.0 and 10.0 mM concentrations at room temperature.  $H_2SO_4$  or  $HNO_3$  were added to the solutions immediately before treatment to adjust pH to 3.5. These solutions were then stored in the dark for 0, 24, 48, and 96 h at room temperature. The samples were rinsed three times with distilled water after treatment and before each assay [29].

### *MDA assay*

For the MDA assay, the Health and Packer [30] method was used. 200 mg of the lichen thalli were homogenized in 2 mL of 0.1% trichloroacetic acid (TCA) and centrifuged at 10000 g for 20 min. To 1 mL of supernatant, 1 mL of 20% TCA containing 0.5 mL of

thiobarbituric acid and 0.001 mL of butylated hydroxyl-toluene BHT at 4% solution in ethanol were added. The mixture was heated at 95 °C for 30 min and centrifuged at 10000 g for 15 min, the supernatant absorbance was read at the wavelength 532 nm and corrected at 600 nm. MDA was calculated by multiplying by the coefficient of  $155 \text{ mM}^{-1}\text{cm}^{-1}$ ; the results were expressed in  $\mu\text{mol g}^{-1}$  FW.

### ***Integrity of chlorophyll assay***

According to Ronen and Galun [31], the integrity of chlorophyll was calculated. About 20 mg of lichen thalli were extracted in 3 mL of dimethylsulfoxide (DMSO) in dark at 65 °C for 40 min. The optical densities at 435 nm and 415 nm for the extract were read, and the ratio  $\text{OD}_{435}/\text{OD}_{415}$  was calculated to assess the degree of chlorophyll degradation. Ronen and Galun (1984) estimate that a ratio between 1.4 and 1.45 was calculated in lichens in the case of minimal chlorophyll degradation into phaeopigments.

### ***Proline assay***

The method used for the determination of proline is that of Troll and Linsley [32], 100 mg of fresh lichen thalli were extracted in 2 mL of 40 % methanol at 85 °C for 60 min. After cooling, 1 mL of acetic acid and 1 mL of a mixture containing (120 mL of distilled water, 300 mL of acetic acid and 80 mL of acid orthophosphoric acid and 25 mg of ninhydrin) were added to 1 mL of the extract. The solution was brought to the boil for 30 min, it gradually turns red. After cooling, 5 mL of toluene was added; the upper phase which contains the proline was recovered and dehydrated by the addition of disodium sulphate. The optical density was determined by a spectrophotometer at a wavelength of 528 nm and the calibration curve was established by different concentrations of proline from a stock solution of 2 mg/100 mL of 40% methanol. Proline contents were determined using equation established by known concentrations of proline ( $y=28.0x$ ,  $R^2=0.9911$ ); the results were expressed in  $\mu\text{g g}^{-1}$  FW.

### ***Soluble sugars assay***

The soluble sugar contents were quantified by the method of Dubois et al. [33]. 100 mg of fresh lichen material were extracted in 3 mL of 85% ethanol for 48 h in the dark, then filtered and recovered with 20 mL of distilled water, 1 mL of 5 % phenol and 5 mL of 1.8 N sulfonic acid were added to 1 mL of the filtrate. After incubation for 15-20 min in a water bath adjusted to 30 °C, the optical densities were determined at the wavelength 490 nm. The calibration curve was established by glucose at different concentrations prepared from a stock solution of 250 mg/L. Soluble sugar contents ( $\mu\text{g.g}^{-1}$  FW) were calculated using equation established by known concentrations of glucose ( $y=0.731x + 0.001$ ,  $R^2=0.999$ ).

### ***Statistical analysis***


Three repetitions were performed at each concentration, so we can calculate the standard deviation (SD). The statistical study was performed using the ORIGIN 6.0 system using the test univariate variance (one way ANOVA). For the study, the results were expressed as mean $\pm$ SD. The difference was considered to be not significant when  $p>0.05^{(NS)}$ , significant when  $0.01<p<0.05$  (\*), highly significant when  $0.001<p<0.01$  (\*\*) and very highly significant when  $p<0.001$  (\*\*\*).

Correlation matrices between NaF, Pb(NO<sub>3</sub>)<sub>2</sub> and different studied parameters were analyzed by STATISTICA Version 10 software.

## RESULTS AND DISCUSSION


### MDA accumulation

Results of the MDA contents in the thalli of *X. parietina* treated by different concentrations of NaF and Pb(NO<sub>3</sub>)<sub>2</sub> during 0, 24, 48 and 96 h are presented in Fig. 1. Correlation matrices between NaF / MDA, Pb(NO<sub>3</sub>)<sub>2</sub> / MDA are presented in Fig. 2.



**Fig. 1.** MDA accumulation in *X. parietina* treated with different concentrations of (A): NaF, (B): Pb(NO<sub>3</sub>)<sub>2</sub>

According to Fig. 1, no significant increase in the MDA content was noted as a function of the concentrations of NaF ( $p=0.179^{NS}$ ) and Pb(NO<sub>3</sub>)<sub>2</sub> ( $p=0.109^{NS}$ ). However, depending on exposure time, a significant increase of MDA content was noted with treatment by NaF ( $=0.005^{**}$ ) and Pb(NO<sub>3</sub>)<sub>2</sub> ( $p=0.014^*$ ).




**Fig. 2.** Correlation matrices between NaF / MDA (A), Pb(NO<sub>3</sub>)<sub>2</sub>/MDA (B).  
(A):  $r=0.773$ ,  $p=0.000712^{***}$ , (B):  $r=0.865$ ,  $p=0.000031^{***}$

From the data presented in Fig. 2, the statistical analysis results shows a significant positive correlation between MDA (Fig. 2 A) and NaF concentration, and between MDA and  $Pb(NO_3)_2$  concentration (Fig. 2 B).

During lipid peroxidation, a wide variety of aldehydes are formed, among these aldehydes is MDA. MDA is used as an interesting biomarker and diagnostic for lipid oxidative damage under drought stress [34, 35]. MDA is also accumulated in plants under heavy metal stress such as mercury [36]. Compared to the control test, our results show that the exposure of *X. parietina* to NaF and  $Pb(NO_3)_2$  solutions caused a significant increase in MDA content ( $p < 0.05^*$ ) where we found a positive correlation between MDA contents in *X. parietina* and increasing concentrations of NaF and  $Pb(NO_3)_2$  ( $r = 0.773$ ,  $p = 0.000712^{***}$  and  $r = 0.865$ ,  $p = 0.000031^{***}$  respectively). The same results were obtained by Dzubaj et al. [37] and Pisani et al. [38] who showed that *X. parietina* reacts against fluorine, boron and lead-induced stress by increasing of the MDA content. Our results are in agreement with those of El-Shora et al. [39] which showed that lead stress increased MDA contents in the treated plants, and those of Alsherif et al. [40] which indicated that heavy metal contamination resulted in significant increases in MDA in plants. Also, our results are in the same line with those obtained by Fan et al. [41] who show that MDA content increased in *Festuca arundinacea* Schreb after high concentration of fluorine treatment. Kacienė et al. [42] showed that oxidative stress induced by stress factors of different origin-ozone, ultraviolet (UV)-B radiation, drought, cadmium and copper, causes the increase in the content of MDA in barley. Likewise, Gutiérrez-Martínez et al. [43] noticed that the MDA content increases with increasing concentrations of cadmium accumulated in the leaves and roots of *Phaseolus vulgaris* plants under cadmium stress.


### Chlorophyll integrity variations

Variations of the  $OD_{435}/OD_{415}$  ratio in *X. parietina* are presented in Fig. 3. Correlation matrices between NaF/ $OD_{435}/OD_{415}$  ratio,  $Pb(NO_3)_2/OD_{435}/OD_{415}$  ratio are presented in Fig. 4.



**Fig. 3.** Variations of the  $OD_{435}/OD_{415}$  ratio in *X. parietina* treated with different concentrations of (A): NaF, (B):  $Pb(NO_3)_2$

Depending on exposure time, Fig. 3 (A) shows that the variations of  $OD_{435}/OD_{415}$  ratio in the thalli treated with the different concentrations of NaF are significant ( $p = 0.009^{**}$ ), but depending of concentrations they are not significant ( $p=0.422^{NS}$ ). A slight decrease in  $OD_{435}/OD_{415}$  ratio in thalli treated with the different concentrations of  $Pb(NO_3)_2$  is recorded after 24 h of treatment ( $p>0.05^{NS}$ ), while between 24 h and 96 of treatment, a significant decrease in this ratio was noticed ( $p=0.036^*$ ).




**Fig. 4.** Correlation matrices between NaF/ $OD_{435}/OD_{415}$  ratio (A),  $Pb(NO_3)_2/OD_{435}/OD_{415}$  ratio (B). (A):  $r = -0.737$ ,  $p = 0.0011^{**}$ , (B):  $r = -0.926$ ,  $p < 0.0001^{***}$

From the data presented in Fig. 4, the statistical analysis results shows a significant negative correlation between  $OD_{435}/OD_{415}$  ratio and exposure time of NaF (Fig. 4 A) and between  $OD_{435}/OD_{415}$  ratio and exposure time of  $Pb(NO_3)_2$  (Fig. 4 B).

The most commonly used metric to quantify chlorophyll degradation is the ratio of optical density of chlorophyll samples read at 435 and 415nm. A ratio of 1.4 informs about chlorophyll integrity, any reduction in this value indicates the degradation of chlorophyll to provide stress to the organism [44, 45]. Ours results show that a ratio of 1.4 was obtained in lichen thalli treated with distilled water (control), but for those treated with different concentrations of NaF and  $Pb(NO_3)_2$ , a decrease in this ratio was noted. According to this results, we found that the decrease of the  $OD_{435}/OD_{415}$  ratio is in correlation with increasing exposure time of *X. parietina* to NaF and  $Pb(NO_3)_2$  ( $r = -0.737$ ,  $p = 0.0011^{**}$  and  $r = -0.926$ ,  $p < 0.0001^{***}$  respectively). Our results are in agreement with those obtained by Shukla and Upreti [46], who reported that  $OD_{435}/OD_{415}$  ratio values decreased with the increase in the amount of Cu, Pb and Zn in the lichen *Pyxine subcinerea* Stirton and of Bajpai et al. [47]; Sharma and Singh [48] and Chetia et al. [49] who indicate that chlorosis increases in lichens under heavy metals stress and with those obtained by Panda [50], who reported that chlorosis is one of the symptoms of fluoride toxicity in plants.

### **Proline accumulation**


Treatment with increasing exposure time to increasing concentrations of NaF and  $Pb(NO_3)_2$  caused a general increasing accumulation of proline in *X. parietina* (Fig. 5). Correlation matrices between NaF/proline,  $Pb(NO_3)_2$ /proline are presented in Fig. 6.



**Fig. 5.** Accumulation of proline in *X. parietina* treated with different concentrations of (A): NaF, (B):  $Pb(NO_3)_2$

According to the Fig. 5 (A), it was noticed that the accumulation of proline in thalli increases significantly with increasing NaF concentrations ( $p=0.0052^{**}$ ) but not significantly with increasing exposure time ( $p=0.177^{NS}$ ).

According to Fig. 5 (B), no significant accumulation of proline in the thalli treated with the 0.5, 1 and 5 mM concentrations of  $Pb(NO_3)_2$  was noted ( $p=0.08^{NS}$ ), however with the high concentration of  $Pb(NO_3)_2$  (10 mM), proline accumulation was significant ( $p=0.016^*$ ). Depending on exposure time, significant accumulation of proline ( $p=0.013^*$ ) after 96 h of treatment with all concentrations of  $Pb(NO_3)_2$  was noted. Therefore, it can be concluded that the accumulation of proline varies much more with time than with concentration.



**Fig. 6.** Correlation matrices between NaF/proline (A),  $Pb(NO_3)_2$ /proline (B). (A):  $r=0.783$ ,  $p=0.00032^{***}$ , (B):  $r=0.811$ ,  $p=0.00013^{***}$


From the data presented in Fig. 6, the statistical analysis results shows a significant positive correlation between proline and exposure time of NaF (Fig. 6 A), and between proline (Fig. 6 B) and exposure time of  $Pb(NO_3)_2$ .

Proline is part of a general adaptive syndrome to adverse environmental conditions [51, 52]. The accumulation of proline can be considered as a biomarker of stress [34] which varies depending on the plant species. Plants react against stress by accumulating

proline to protect the structure of its macromolecules [53]. Amri and Layachi [54] reported that exogenous application of proline on a Faba bean (*Vicia faba*) plant cultivated under cadmium stress helped the plant recover from the cadmium stress-induced physiological changes. The determination of proline in lichens is a detective method of the various possible stress phenomena. Our results show a significant increase in proline content in *X. parietina* correlating with increasing exposure time to NaF and  $\text{Pb}(\text{NO}_3)_2$  ( $r=0.783$ ,  $p=0.00032^{***}$  and  $r=0.811$ ,  $p=0.00013^{***}$  respectively), the same result was obtained by Li et al. [55] who reported that proline contents increased in maize varieties under Cd stress. Likewise, our results are similar to those obtained by Koleva et al. [56], which were noted that *Phaseolus vulgaris* seedlings under cadmium-induced stress exhibited an increased level of proline. Several other studies indicate also that proline increases under the action of other types of stress: salt stress [53], water limitation [57], changing climate conditions [52], UV radiation [58], heat stress tolerance [59] and nutrient deficiency [28].


### *Soluble sugars accumulation*

The effect of different concentrations of NaF and  $\text{Pb}(\text{NO}_3)_2$  on the content of soluble sugars in *X. parietina* are shown in Fig. 7. Correlation matrices between NaF/soluble sugars,  $\text{Pb}(\text{NO}_3)_2$ /soluble sugars are presented in Fig. 8.



**Fig. 7.** variations of soluble sugar contents in *X. parietina* treated with different concentrations of (A): NaF, (B):  $\text{Pb}(\text{NO}_3)_2$

According to the results presented in Fig. 7, the soluble sugar contents generally increase in thalli treated with different concentrations of NaF and  $\text{Pb}(\text{NO}_3)_2$ , but these contents decrease slightly after 96 h of  $\text{Pb}(\text{NO}_3)_2$  treatment with 5 mM and 10 mM concentrations and with all concentrations of NaF (0.5 mM, 1 mM, 5 mM, and 10 mM). Fig. 7 (A) shows a significant increase in soluble sugar contents in thalli treated with all concentrations of NaF ( $p=0.01^*$ ), but depending of exposure time, the variations in soluble sugar contents were not significant ( $p=0.23^{NS}$ ). Unlike NaF, Fig. 7 (B) shows no significant increase in soluble sugar contents depending of concentrations ( $p=0.238^{NS}$ ), and a significant increase depending of exposure time ( $p=0.010^*$ ).



**Fig. 8.** Correlation matrices between NaF/soluble sugars (A), Pb(NO<sub>3</sub>)<sub>2</sub>/soluble sugars (B). (A):  $r=0.678$ ,  $p=0.0010^{***}$ , (B):  $r=0.780$ ,  $p=0.00036^{***}$

From the data presented in Fig. 8, the statistical analysis results shows a significant positive correlation between soluble sugars (Fig. 8 A) and NaF concentration, and between soluble sugars contents (Fig. 8 B) and exposure time of Pb(NO<sub>3</sub>)<sub>2</sub>.

Like proline, soluble sugars are part of the adaptation strategies used by plants to combat and tolerate stressful conditions [27, 34, 60]. The results obtained show a significant increase in the content of soluble sugars correlating with increasing concentration of NaF ( $r=0.678$ ,  $p=0.0010^{***}$ ) and correlating with increasing exposure time of *X. parietina* to Pb(NO<sub>3</sub>)<sub>2</sub> ( $r=0.780$ ,  $p=0.00036^{***}$ ); these results are in agreement with the results obtained by Gandonou et al. [61] who showed that soluble sugars accumulate in two sugarcane cultivars under salt stress. According to Abbaspour et al. [62], the increase of the salinity stress resulted in the increasing concentration of soluble sugars in three pistachio cultivars. Also, several other studies showed that drought stress increased the contents of soluble sugars in the leaves of soybean seedlings [63] and in *Sophora davidii* (Franch.) [64]. Our results are also concomitant with those obtained by Aldoobie and Beltagi [65], who reported that contents of total soluble sugars increased in common bean (*Phaseolus vulgaris* L. cv. Nebraska) plants in response to lead, cadmium and nickel stress.

According to Ahmad et al. [66], plants accumulate soluble sugars as a defense mechanism against stressful conditions caused by drought and water scarcity, varying temperature from minimal to maximum level, and accumulation of salt and heavy metals.

Our results show that chlorosis, lipid peroxidation, accumulation of proline and soluble sugars in *X. parietina* under Pb(NO<sub>3</sub>)<sub>2</sub> treatment are more intense than under NaF treatment. Based on our statistical analyses, it was concluded that lead is significantly more toxic than fluoride ( $p=0.02^*$ ).

## CONCLUSION

The results of the present study showed a significant increase of MDA content in *X. parietina* correlating with increasing concentrations of NaF and Pb(NO<sub>3</sub>)<sub>2</sub>, and a significant increase of OD<sub>435</sub>/OD<sub>415</sub> ratio and proline accumulation correlating with increasing exposure time of NaF and Pb(NO<sub>3</sub>)<sub>2</sub>, results showed also that soluble sugar contents increased correlating with increasing concentrations of NaF and correlating with increasing exposure time of Pb(NO<sub>3</sub>)<sub>2</sub>. Despite the fact that lead was more toxic than

fluoride, *X. parietina* has a very high sensitivity to fluoride, we were able to draw the conclusion that fluoride must be classified among the most toxic air pollutants, and therefore to open the field to other works to study and compare the toxicity of fluorine with that of heavy metals on the various other ecosystems.

**Acknowledgement.** The author thanks the collaborators for their contributions during experimental studies.

**Conflict of Interest.** The author declared that there is no conflict of interest.

**Authorship Contributions.** Concept: O.B., N.B., E.L., Design: O.B., N.B., E.L., Data Collection or Processing: O.B., N.B., E.L., Analysis or Interpretation: O.B., N.B., E.L., Literature Search: O.B., N.B., E.L., Writing: O.B., N.B., E.L.

**Financial Disclosure.** This research received no grant from any funding agency/sector.

## REFERENCES

- [1] Nash, III T. (2008): Lichen Biology (2nd Ed.). Cambridge: Cambridge, University Press. DOI: <https://doi.org/10.1017/CBO9780511790478>.
- [2] Benítez, A., Medina, J., Vásquez, C., Loaiza, T., Luzuriaga, Y., Calva, J. (2019): Lichens and Bromeliads as Bioindicators of Heavy Metal Deposition in Ecuador. Lichen Diversity and Biomonitoring 11(2), 28. DOI: <https://doi.org/10.3390/d11020028>.
- [3] Mohamed, E., Mohamed, L., Abdelhay, E.G. (2020): Using calcicolous and corticolous lichens to assess lead and cadmium air pollution of the Moroccan Atlantic Coast Safi-Essaouira. Polish Journal of Environmental Studies 29(1): 779-787. DOI: <https://doi.org/10.15244/pjoes/102629>.
- [4] Quijano-Abril, M.A., Ramirez-Ospina, D.M., Domínguez-Rave, M.I., Londoño-Valencia, J. (2021): Lichens as biosensors for the evaluation of urban and sub-urban air pollution in a tropical mountain valley, Rionegro, Antioquia. Revista Bionatura 6(1): 1501-1509. DOI: <https://doi.org/10.21931/RB/2021.06.01.10>.
- [5] Caggiano, R., Trippetta, S., Sabia, S. (2015): Assessment of atmospheric trace element concentrations by lichen-bag near an oil/gas pre-treatment plant in the Agri Valley (southern Italy). Natural Hazards and Earth System Sciences 15(2): 325-333. DOI: <https://doi.org/10.5194/nhess-15-325-2015>.
- [6] Darnajoux, R., Lutzoni, F., Miadlikowska, J., Bellenger, J.P. (2015): Determination of elemental baseline using peltigeralean lichens from Northeastern Canada (Québec): Initial data collection for long term monitoring of the impact of global climate change on boreal and subarctic area in Canada. Science of the Total Environment 533: 1-7. DOI: <https://doi.org/10.1016/j.scitotenv.2015.06.030>
- [7] Demková, L., Bobul'ská, L., Árvay, J., Jezný, T., Ducsay, L. (2017): Biomonitoring of heavy metals contamination by mosses and lichens around Slovinky tailing pond (Slovakia). Journal of Environmental Science and Health, Part A, Toxic/Hazardous Substances and Environmental Engineering 52(1): 30-36. DOI: <https://doi.org/10.1080/10934529.2016.1221220>.
- [8] Abas, A. (2021): A systematic review on biomonitoring using lichen as the biological indicator: A decade of practices, progress and challenges. Ecological Indicators 121: 107-197. DOI: <https://doi.org/10.1016/j.ecolind.2020.107197>.
- [9] Tarawneh, A.H., Salamon, I., Altarawneh, R., Mitra, J.(2021): Assessment of Lichens as Biomonitorers of Heavy Metal Pollution in Selected Mining Area, Slovakia. Pakistan Journal

- of Analytical and Environmental Chemistry 22(1): 53-59. DOI: <https://doi.org/10.21743/pjaec/2021.06.07>.
- [10] Węgrzyn, M., Wietrzyk, P., Lisowska, M., Klimek, B., Nicia, P. (2016): What influences heavy metals accumulation in arctic lichen *Cetrariella delisei* in Svalbard? Polar Science 10(4): 532-540. DOI: <https://doi.org/10.1016/j.polar.2016.10.002>.
- [11] Winkler, A., Caricchi, C., Guidotti, M., Owczarek, M., Macri, P., Nazzari, M., Amoroso, A., Di Giosa, A., Listrani, S. (2019): Combined magnetic, chemical and morphoscopic analyses on lichens from a complex anthropic context in Rome. Italy, Science of The Total Environment 690: 1355-1368. DOI: <https://doi.org/10.1016/j.scitotenv.2019.06.526>.
- [12] Rola, K. (2020): Insight into the pattern of heavy-metal accumulation in lichen thalli. Journal of Trace Elements in Medicine and Biology 61: 126512. DOI: <https://doi.org/10.1016/j.jtemb.2020.126512>.
- [13] Vannini, A., Tedesco, R., Loppi, S., Di Cecco, V., Di Martino, L., Nascimbene, J., Dallo, F., Barbante, C. (2021): Lichens as monitors of the atmospheric deposition of potentially toxic elements in high elevation Mediterranean ecosystems. Science of The Total Environment 798: 149369. DOI: <https://doi.org/10.1016/j.scitotenv.2021.149369>.
- [14] Dévéhat, F., Thüs, H., Abasq, M.L., Delmail, D., Boustie, J. (2014): Oxidative Stress Regulation in Lichens and Its Relevance for Survival in Coastal Habitats. Advances in Botanical Research 71: 467-503. DOI: <https://doi.org/10.1016/B978-0-12-408062-1.00016-0>.
- [15] Expósito, J.R., Barreno, E., Catalá, M. (2022): 18 - Role of NO in lichens. In: Singh VP, Singh S., Tripathi D.K., Romero-Puertas M.C., Sandalio M.L. (eds), *Nitric Oxide in Plant Biology*. Academic Press, pp. 407-429. DOI: <https://doi.org/10.1016/B978-0-12-818797-5.00027-3>.
- [16] Škvorová, Z., Černajová, I., Steinová, J., Peksa, O., Moya, P., Škaloud, P. (2022): Promiscuity in Lichens Follows Clear Rules: Partner Switching in *Cladonia* Is Regulated by Climatic Factors and Soil Chemistry. *Frontiers in Microbiology* 12. DOI: <https://doi.org/10.3389/fmicb.2021.781585>.
- [17] Beckett, R., Minibayeva, F., Solhaug, K., Roach, T. (2021): Photoprotection in lichens: Adaptations of photobionts to high light. *The Lichenologist* 53(1): 21-33. DOI: <https://doi.org/10.1017/S0024282920000535>.
- [18] Chowaniec, K. and Rola, K. (2022): Evaluation of the importance of ionic and osmotic components of salt stress on the photosynthetic efficiency of epiphytic lichens. *Physiology and Molecular Biology of Plants* 28: 107–121. DOI: <https://doi.org/10.1007/s12298-022-01134-2>.
- [19] Kraft, M., Scheidegger, C., Werth, S. (2022): Stressed out: the effects of heat stress and parasitism on gene expression of the lichen-forming fungus *Lobaria pulmonaria*. *The Lichenologist* 54: 71–83. DOI: <https://doi.org/10.1017/S0024282921000463>.
- [20] Hauck, M., Willenbruch, K., Leuschner, C. (2009): Lichen Substances Prevent Lichens from Nutrient Deficiency. *Journal of Chemical Ecology* 35: 71–73. DOI: <https://doi.org/10.1007/s10886-008-9584-2>.
- [21] Kranner, I., Beckett, R., Hochman, A., Nash, III.T. (2009): Desiccation-Tolerance in Lichens: A Review. *The Bryologist* 111: 576-593. DOI: <https://doi.org/10.1639/0007-2745-111.4.576>.
- [22] Armstrong, R.A. (2017): Adaptation of Lichens to Extreme Conditions. In: Shukla V, Kumar S, Kumar N. (eds.). *Plant Adaptation Strategies in Changing Environment*. Springer, Singapore, pp. 1-27. DOI: [https://doi.org/10.1007/978-981-10-6744-0\\_1](https://doi.org/10.1007/978-981-10-6744-0_1).
- [23] Karakoti, N., Bajpai, R., Upreti, D.K., Mishra, G.K., Srivastava, A., Nayaka, S. (2014): Effect of metal content on chlorophyll fluorescence and chlorophyll degradation in lichen *Pyxine cocolos* (Sw.) Nyl.: a case study from Uttar Pradesh, India. *Environmental Earth Sciences* 71(5). DOI: <https://doi.org/10.1007/s12665-013-2623-5>.
- [24] Sujetovienė, G. (2015): Monitoring Lichen as Indicators of Atmospheric Quality. In: Upreti D, Divakar P, Shukla V, Bajpai R. (eds), *Recent Advances in Lichenology*. Vol. 1. Modern

- Methods and Approaches in Biomonitoring and Bioprospection. Springer, New Delhi, pp. 87-118. DOI: [https://doi.org/10.1007/978-81-322-2181-4\\_4](https://doi.org/10.1007/978-81-322-2181-4_4).
- [25] Paoli, L., Munzi, S., Guttova, A., Senko, D., Sardella, G., Loppi, S. (2015) : Lichens as suitable indicators of the biological effects of atmospheric pollutants around a municipal solid waste incinerator (S Italy). *Ecological Indicators* 52. DOI: <https://doi.org/10.1016/j.ecolind.2014.12.018>.
- [26] Sujetovienė, G., Smilgaitis, P., Dagiliūtė, R., Žaltauskaitė, J. (2019): Metal accumulation and physiological response of the lichens transplanted near a landfill in central Lithuania. *Waste Management* 85: 60-65. DOI: <https://doi.org/10.1016/j.wasman.2018.12.017>.
- [27] Khan, N., Ali, S., Zandi, P., Mehmood, A., Ullah, S., Ikram, M., Ismail, I., Shahid, M., Babar, Md. (2020): Role of sugars, amino acids and organic acids in improving plant abiotic stress tolerance. *Pakistan Journal of Botany* 52(2). DOI: [https://doi.org/10.30848/PJB2020-2\(24\)](https://doi.org/10.30848/PJB2020-2(24)).
- [28] Mundada, P.S., Jadhav, S.V., Salunkhe, S.S., Gurme, S.T., Umdale, S.D., Nikam, T.D., Ahire, M.L. (2021): Plant Performance and Defensive Role of Proline Under Environmental Stress. In: Husen A. (ed.), *Plant Performance Under Environmental Stress*. Springer; Berlin/Heidelberg, pp. 201-223. DOI: [https://doi.org/10.1007/978-3-030-78521-5\\_8](https://doi.org/10.1007/978-3-030-78521-5_8).
- [29] Carreras, H.A. and Pignata, M.L. (2007): Effects of the heavy metals Cu<sup>2+</sup>, Ni<sup>2+</sup>, Pb<sup>2+</sup>, and Zn<sup>2+</sup> on some physiological parameters of the lichen *Usnea amblyoclada*. *Ecotoxicology and Environmental Safety* 67(1):59-66. DOI: <https://doi.org/10.1016/j.ecoenv>.
- [30] Heath, R.L. and Packer, L. (1968): Photoperoxidation in isolated chloroplasts: I. Kinetics and stoichiometry of fatty acid peroxidation. *Archives of Biochemistry and Biophysics* 125(1): 189-198. DOI: [https://doi.org/10.1016/0003-9861\(68\)90654-1](https://doi.org/10.1016/0003-9861(68)90654-1).
- [31] Ronen, R. and Galun, M. (1984): Pigment extraction from lichens with dimethyl sulfoxide (DM50) and estimation of chlorophyll degradation. *Environmental and Experimental Botany* 24: 239-245. DOI: [http://dx.doi.org/10.1016/0098-8472\(84\)90004-2](http://dx.doi.org/10.1016/0098-8472(84)90004-2).
- [32] Troll, W. and Lindsley, J. (1955): A photometric method for the determination of proline. *Journal of Biological Chemistry* 215: 655-60. DOI: [https://doi.org/10.1016/S0021-9258\(18\)65988-5](https://doi.org/10.1016/S0021-9258(18)65988-5).
- [33] Dubois, M, Gilles, KA, Hamilton, JK, Rebers, PA, Smith, F. (1956): *Analytical Chemistry* 28, 350-356.
- [34] Amine-Khodja, I.R., Boscari, A., Riah, N., Kechid, M., Maougal, R.T., Belbekri, N., Djekoun, A. (2022): Impact of Two Strains of *Rhizobium leguminosarum* on the Adaptation to Terminal Water Deficit of Two Cultivars *Vicia faba*. *Plants* 11(4): 515. DOI: <https://doi.org/10.3390/plants11040515>.
- [35] Toto, A., Wild, P., Graille, M., Turcu, V., Crézé, C., Hemmendinger, M., Sauvain, J.J., Bergamaschi, E., Canu, I.G., Hopf, N.B. (2022): Urinary Malondialdehyde (MDA) Concentrations in the General Population-A Systematic Literature Review and Meta-Analysis. *Toxics* 10(4): 160. DOI: <https://doi.org/10.3390/toxics10040160>.
- [36] Singh, H., Kumar, D., Soni, V. (2020): Copper and mercury induced oxidative stresses and antioxidant responses of *Spirodela polyrhiza* (L.) Schleid. *Biochemistry and Biophysics Reports* 23: 100781. DOI: <https://doi.org/10.1016/j.bbrep.2020.100781>.
- [37] Dzubaj, A., Backor, M., Tomko, J., Péli, E. and Tuba, Z. (2008): Tolerance of the lichen *Xanthoria parietina* (L.) Th. Fr. to metal stress. *Ecotoxicology and Environmental Safety* 70(2): 319-26. DOI: <https://doi.org/10.1016/j.ecoenv.2007.04.002>.
- [38] Pisani, T., Munzi, S., Paoli, L., Backor, M., Loppi, S. (2009): Physiological effects of a geothermal element: boron excess in the epiphytic lichen *Xanthoria parietina* (L.) TH. FR. *Chemosphere* 76(7): 921-6. DOI: <https://doi.org/10.1016/j.chemosphere.2009.04.058>.
- [39] El-Shora, H., Massoud, G., Gad, D. (2021): Activation of metabolic pathways associated with phenolic biosynthesis in Garden Cress leaves under lead stress. *Project: Plant metabolites under stress*. 11: 128-140. DOI: <https://doi.org/10.33887/rjpbcs/2020.11.6.16>.

- [40] Alsherif, E.A., Al-Shaikh, T.M. and AbdElgawad, H. (2022): Heavy Metal Effects on Biodiversity and Stress Responses of Plants Inhabiting Contaminated Soil in Khulais. Saudi Arabia. *Biology* 11(2): 164. DOI: <https://doi.org/10.3390/biology11020164>.
- [41] Fan, F., Chen, K., Xu, J., Abm, K., Chen, Y., Chen, L., and, Yan, X. 2022. Physiological effects induced by aluminium and fluoride stress in tall fescue (*Festuca arundinacea* Schreb). *Ecotoxicology and Environmental Safety* 231: 113192. DOI: <https://doi.org/10.1016/j.ecoenv.2022.113192>
- [42] Kacienė, G., Žaltauskaitė, J., Milčė, E., Juknys, R. (2015): Role of oxidative stress on growth responses of spring barley exposed to different environmental stressors. *Journal of Plant Ecology* 8 (6): 605–616. DOI: <https://doi.org/10.1093/jpe/rtv026>.
- [43] Gutiérrez-Martínez, P.B., Torres-Morán, M.I., Romero-Puertas, M.C., Casas-Solís, J., Zarazúa-Villaseñor, P., Sandoval-Pinto, E., Ramírez-Hernández, B.C. (2020): Assessment of antioxidant enzymes in leaves and roots of *Phaseolus vulgaris* plants under cadmium stress. *Biocencia* 22 (2): 110-118. DOI: <https://doi.org/10.18633/biocencia.v22i2.1252>.
- [44] Munzi, S., Pirintsos, S.A., Loppi, S. (2009): Chlorophyll degradation and inhibition of polyamine biosynthesis in the lichen *Xanthoria parietina* under nitrogen stress. *Ecotoxicology and Environment Safety* 72: 281-285. DOI: <https://doi.org/10.1016/j.ecoenv.2008.04.013>
- [45] Bajpai, R., Upreti, D.K., Nayakaa, S., Kumarib, B. (2010): Biodiversity, bioaccumulation and physiological changes in lichens growing in the vicinity of coal-based thermal power plant of Raebareli district, north India. *Journal of Hazard Materials* 174: 429-436. DOI: <http://dx.doi.org/10.1016/j.jhazmat.2009.09.071>.
- [46] Shukla, V. and Upreti, D. (2008): Effect of metallic pollutants on the physiology of lichen, *Pyxine subcinerea* Stirton in Garhwal Himalayas. *Environmental Monitoring and Assessment* 141: 237-243. DOI: <https://doi.org/10.1007/S10661-007-9891-Z>.
- [47] Bajpai, R., Shukla, V., Singh, N., Rana, T.S., Upeti, D.K. (2015): Physiological and genetic effects of chromium (+VI) on toxitolerant lichen species *Pyxine cocoes*. *Environmental Science and Pollution Research* 22(5): 3727-3738. DOI: <http://dx.doi.org/10.1007/s11356-014-3622-0>.
- [48] Sharma, R. and Singh, R. (2016): Air pollution biomonitoring in Delhi city by using lichen transplant technique. *Cryptogam Biodiversity and Assessment* 1(1): 55 – 63. DOI: <https://doi.org/10.21756/cba.v1i1.11018>.
- [49] Chetia, J., Gogoi, N., Gogoi, R., Yasmin, F. (2021): Impact of heavy metals on physiological health of lichens growing in differently polluted areas of central Assam, North East India. *Plant Physiology Reports* 26: 210-219. DOI: <https://doi.org/10.1007/s40502-021-00575-3>.
- [50] Panda, D. (2015): Fluoride toxicity stress: physiological and biochemical consequences on plants. *International journal of bio-resource, environment and agricultural sciences* 1(1): 70-84.
- [51] Liang, X., Zhang, L., Natarajan, S.K., Becker, D.F. (2013): Proline Mechanisms of Stress Survival. *Antioxidants and Redox Signaling* 19(9). DOI: <https://doi.org/10.1089/ars.2012.5074>.
- [52] Ghosh, U.K., Islam, M.N., Siddiqui, M.N., Cao, X., Khan, M.A.R. (2022): Proline, a multifaceted signalling molecule in plant responses to abiotic stress: understanding the physiological mechanisms. *Plant Biology* 24(2):227-239. DOI: <https://doi.org/10.1111/plb.13363>.
- [53] Alhasnawi, A. (2019): Role of proline in plant stress tolerance: A mini review. *Resurrect Crops* 20(1): 223-229. DOI: <https://doi.org/10.31830/2348-7542.2019.032>.
- [54] Amri, A. and Layachi, N. (2018): Interactive effects of cadmium stress and proline on physiological and biochemical parameters of faba bean plant. *International Journal of Biosciences* 12(5): 88-100. DOI: <http://dx.doi.org/10.12692/ijb/12.5.88-100>.

- [55] Li, C, Cao, Y, Li, T, Guo, M, Ma, X, Zhu, Y, Fan, J. (2022) : Changes in antioxidant system and sucrose metabolism in maize varieties exposed to Cd. *Environmental Science and Pollution Research* 15(12): e0243835. DOI: <https://doi.org/10.1007/s11356-022-20422-8>
- [56] Koleva, L., Umar, A., Yasin, N.A., Shah, A.A., Siddiqui, M.H., Alamri, S., Riaz, L., Raza, A., Javed, T., Shabbir, Z. (2022): Iron Oxide and Silicon Nanoparticles Modulate Mineral Nutrient Homeostasis and Metabolism in Cadmium-Stressed *Phaseolus vulgaris*. *Frontiers in Plant Science* 13: 806781. DOI: <https://doi.org/10.3389/fpls.2022.806781>.
- [57] Bhaskara, G.B., Yang, T.H., Verslues, P.E. (2015): Dynamic proline metabolism: Importance and regulation in water limited environments. *Frontiers in Plant Science* 6, 484. DOI: <https://doi.org/10.3389/fpls.2015.00484>.
- [58] Metwally, S.A., Shoaib, R.M., Hashish, K.I., El-Tayeb, T.A. (2019): In vitro ultraviolet radiation effects on growth, chemical constituents and molecular aspects of *Spathiphyllum* plant. *Bulletin of the National Research Centre* 43(94). DOI: <https://doi.org/10.1186/s42269-019-0126-6>.
- [59] Iqbal, N., Fatma, M., Khan, N., Umar, S. (2019): Regulatory Role of Proline in Heat Stress Tolerance. In: *Plant Signaling Molecules*. Woodhead Publishing, pp. 437-448. DOI: <https://doi.org/10.1016/B978-0-12-816451-8.00027-7>.
- [60] Gangola, M.P. and Ramadoss, B.R. (2018): Chapter 2 - Sugars Play a Critical Role in Abiotic Stress Tolerance in Plants. In: Wani SH. (ed.), *Biochemical, Physiological and Molecular Avenues for Combating Abiotic Stress Tolerance in Plants*, first ed. Academic Press, pp. 17-38. DOI: <https://doi.org/10.1016/B978-0-12-813066-7.00002-4>.
- [61] Gandonou, C.B., Bada, F., Abrini, J., Skali-Senhaji, N. (2011): Free proline, soluble sugars and soluble proteins concentration as affected by salt stress in two sugarcane (*Saccharum sp.*) cultivars differing in their salt tolerance. *International Journal of Biological and Chemical Sciences* 5(6): 2441-2453. DOI: <http://dx.doi.org/10.4314/ijbcs.v5i6.23>.
- [62] Abbaspour, H., Afshari, H., Abdel-Wahhab, M.A. (2012): Influence of salt stress on growth, pigments, soluble sugars and ion accumulation in three pistachio cultivars. *Journal of Medicinal Plants Research* 6(12): 2468-2473. DOI: <https://doi.org/10.5897/JMPR11.1710>.
- [63] Du, Y., Zhao, Q., Chen, L., Yao, X., Zhang, W., Zhang, B., Xie, F. (2020): Effect of drought stress on sugar metabolism in leaves and roots of soybean seedlings. *Plant Physiology and Biochemistry* 146: 1-12. DOI: <https://doi.org/10.1016/j.plaphy.2019.11>.
- [64] Zhao, X., Huang, L.J., Sun, X.F., Zhao, L.L., Wang, P.C. (2022): Transcriptomic and Metabolomic Analyses Reveal Key Metabolites, Pathways and Candidate Genes in *Sophora davidii* (Franch.) Skeels Seedlings Under Drought Stress. *Frontiers in Plant Science* 13. DOI: <https://doi.org/10.3389/fpls.2022.785702>.
- [65] Aldoobie, N.F. and Beltagi, M.S. (2013): Physiological, biochemical and molecular responses of common bean (*Phaseolus vulgaris* L.) plants to heavy metals stress. *African Journal of Biotechnology* 12(29): 4614-4622. DOI: <https://doi.org/10.5897/AJB2013.12387>.
- [66] Ahmad, F., Singh, A., Kamal, A. (2020): Osmoprotective Role of Sugar in Mitigating Abiotic Stress in Plants. In: Roychoudhury A, Tripathi DK. (eds.). *Protective Chemical Agents in the Amelioration of Plant Abiotic Stress*. Biochemical and Molecular Perspectives pp. 53-70. DOI: <https://doi.org/10.1002/9781119552154>.